

HELP GET READY FOR YOUR UPCOMING PROCEDURE

Doptelet[®]
(avatrombopag) tablets

DOPTELET is a prescription medicine used to treat low blood platelet counts in adults with long-lasting (chronic) liver disease who are scheduled to have a medical or dental procedure.

DOPTELET is not used to make platelet counts normal in adults with chronic liver disease. It is not known if DOPTELET is safe and effective in children.

**READY.
SET.
DOPTELET.**

SELECTED IMPORTANT SAFETY INFORMATION

The most common side effects of DOPTELET in people with chronic liver disease are fever, stomach (abdominal) pain, nausea, headache, tiredness, and swelling of the hands or feet. These are not all the possible side effects of DOPTELET.

Please see full Important Safety Information on page 11 and accompanying Full Prescribing Information for DOPTELET (avatrombopag), including Patient Information; also available at Doptelet.com.



Doptelet[®] *(avatrombopag) tablets*

**For adults with chronic liver disease and
a low platelet count before a scheduled
medical or dental procedure**

Get to know DOPTELET

This brochure will help you understand why you need to take DOPTELET before your procedure and what you need to do to complete your treatment.

DOPTELET is a prescription medicine that has been prescribed to you because you have a low platelet count (thrombocytopenia), in addition to chronic liver disease.

You should take DOPTELET just as your healthcare provider prescribed.

SELECTED IMPORTANT SAFETY INFORMATION

The most common side effects of DOPTELET are fever, stomach (abdominal) pain, nausea, headache, tiredness, and swelling of the hands or feet. These are not all the possible side effects of DOPTELET.


Please see full Important Safety Information on page 11 and accompanying Full Prescribing Information for DOPTELET (avatrombopag), including Patient Information; also available at Doptelet.com.



Understanding low platelet count

Platelets are blood cells that work together to form clots that help stop bleeding and repair damage. If you bleed, the platelets rush to the site of the bleeding and form a clot or plug to repair the damage.

If your platelets are low, having more platelets on the day of your procedure may help reduce bleeding. Your healthcare provider prescribed DOPTelet® (avatrombopag) tablets for you with this in mind.



SELECTED IMPORTANT SAFETY INFORMATION

The most common side effects of DOPTelet are fever, stomach (abdominal) pain, nausea, headache, tiredness, and swelling of the hands or feet. These are not all the possible side effects of DOPTelet.

Please see full Important Safety Information on page 11 and accompanying Full Prescribing Information for DOPTelet (avatrombopag), including Patient Information; also available at Doptelet.com.

About DOPTELET® (avatrombopag)

Here are some of the most common questions and answers about DOPTELET. If you have any other questions about DOPTELET or your upcoming procedure, speak to your healthcare provider right away.

Q: Why do I need DOPTELET?

A: DOPTELET can help increase your platelet count before your procedure.

Q: How does DOPTELET work?

A: DOPTELET helps your body by creating more platelets. It does not interfere with the normal way you make platelets, but instead adds more to what you already have.

Q: What are the alternatives to DOPTELET?

A: Up until now, a platelet transfusion has been the most common option for people who needed to raise their platelet count before a procedure; however, with platelet transfusions you need to go to a hospital or hospital-like setting (outpatient center) and it can take a few hours.

With DOPTELET, you just take 5 daily doses at home to begin preparing for your procedure.



Doptelet[®]
(avatrombopag) tablets

Q: How should I take DOPTELET® (avatrombopag)?

A: Starting on the right day is important. You should take DOPTELET just as your healthcare provider prescribed. You will begin taking DOPTELET 10 to 13 days prior to your procedure. Someone from your healthcare team, most likely the specialty pharmacy, will contact you to remind you when you should start taking DOPTELET.

Three things you should know about taking DOPTELET:



ONCE A DAY

Take your dose of DOPTELET once a day.



WITH FOOD

Take DOPTELET every day with food.

5
DAYS

FOR 5 DAYS

Take DOPTELET for 5 days in a row as directed by your healthcare provider.

Other things to remember when taking DOPTELET:

- If you are taking DOPTELET prior to a scheduled medical procedure and you miss a dose, contact your healthcare provider for further dosing instructions.
- Complete all 5 doses of medication.

SELECTED IMPORTANT SAFETY INFORMATION

The most common side effects of DOPTELET are fever, stomach (abdominal) pain, nausea, headache, tiredness, and swelling of the hands or feet. These are not all the possible side effects of DOPTELET.

Please see full Important Safety Information on page 11 and accompanying Full Prescribing Information for DOPTELET (avatrombopag), including Patient Information; also available at Doptelet.com.



If you have any questions about how or when to take DOPTELET® (avatrombopag), call your healthcare provider. You can also call your specialty pharmacy, or Dova 1Source™ at 1-833-DOVA-ONE (1-833-368-2663), and they will help you.



Doptelet®
(avatrombopag) tablets

Q: What does DOPTELET® (avatrombopag) look like?

A: DOPTELET will come in a box, with the tablets in a blister pack. Your healthcare provider will prescribe to you either 40 mg—2 tablets per day or 60 mg—3 tablets per day, based on your platelet count.

DOPTELET 40 mg—2 tablets a day for 5 days



DOPTELET 60 mg—3 tablets a day for 5 days



Q: How do I get DOPTELET?

A: There are several different ways for you to get DOPTELET. One way is for a specialty pharmacy to deliver DOPTELET right to your home or to your doctor's office/clinic. Another way is to get DOPTELET directly from the clinic or hospital pharmacy.

Please see full Important Safety Information on page 11 and accompanying Full Prescribing Information for DOPTELET (avatrombopag), including Patient Information; also available at Doptelet.com.

Dova 1Source is here to help you receive your prescription and provide you with the support you may need. Here's how the process works:



1. Your healthcare provider will send your DOPTelet® (avatrombopag) prescription to Dova 1Source.



2. Dova 1Source will call you to review DOPTelet information with you, and send your prescription to your specialty pharmacy.



3. Your specialty pharmacy will contact you to confirm your delivery details before sending DOPTelet to you.



4. After you receive DOPTelet, Dova 1Source will contact you again to address questions that you may have about taking DOPTelet.



Q: What are the possible side effects of DOPTelet® (avatrombopag)?

A: DOPTelet may cause serious side effects, including blood clots. People with chronic liver disease and people with certain blood clotting conditions may have an increased risk of developing blood clots. Tell your healthcare provider right away if you have signs and symptoms of a blood clot, including swelling, pain or tenderness in your leg; shortness of breath; chest pain; fast heartbeat; or stomach (abdominal) pain or tenderness.

The most common side effects of DOPTelet are fever, stomach (abdominal) pain, nausea, headache, tiredness, and swelling of the hands or feet. These are not all the possible side effects of DOPTelet.

Q: What should I tell my healthcare provider before taking DOPTelet?

A: Before you take DOPTelet, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had a blood clot.
- are pregnant or plan to become pregnant. DOPTelet may harm your unborn baby. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with DOPTelet.
- are breastfeeding or plan to breastfeed. It is not known if DOPTelet passes into your breast milk. Do not breastfeed during your treatment with DOPTelet and for at least 2 weeks after the last dose. Talk to your healthcare provider about the best way to feed your baby during this time.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. DOPTelet may affect the way other medicines work, and other medicines may affect the way DOPTelet works.

Please see full Important Safety Information on page 11 and accompanying Full Prescribing Information for DOPTelet (avatrombopag), including Patient Information; also available at Doptelet.com.

What is DOPTelet® (avatrombopag)?

DOPTelet is a prescription medicine used to treat low blood platelet counts in adults with long-lasting (chronic) liver disease who are scheduled to have a medical or dental procedure.

DOPTelet is not used to make platelet counts normal in adults with chronic liver disease. It is not known if DOPTelet is safe and effective in children.

IMPORTANT SAFETY INFORMATION FOR DOPTelet

What are the possible side effects of DOPTelet?

- DOPTelet may cause serious side effects, including blood clots. People with chronic liver disease and people with certain blood clotting conditions may have an increased risk of developing blood clots. Tell your healthcare provider right away if you have signs and symptoms of a blood clot, including swelling, pain or tenderness in your leg; shortness of breath; chest pain; fast heartbeat; or stomach (abdominal) pain or tenderness.
- The most common side effects of DOPTelet are fever, stomach (abdominal) pain, nausea, headache, tiredness, and swelling of the hands or feet. These are not all the possible side effects of DOPTelet.

Before you take DOPTelet, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had a blood clot.
- are pregnant or plan to become pregnant. DOPTelet may harm your unborn baby. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with DOPTelet.
- are breastfeeding or plan to breastfeed. It is not known if DOPTelet passes into your breast milk. Do not breastfeed during your treatment with DOPTelet and for at least 2 weeks after the last dose. Talk to your healthcare provider about the best way to feed your baby during this time.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

For product information and adverse event reports, please contact:

Dova Medical Information Call Center

Phone: 1-844-506-3682

E-mail: medinfo@dova.com

Reference: DOPTelet [package insert].
Durham, NC: Dova Pharmaceuticals, Inc; 2019.



Doptelet®
(avatrombopag) tablets

Dosing Reminder

Take DOPTelet® (avatrombopag) with food, once a day, for 5 days in a row. When someone from the specialty pharmacy contacts you, they will remind you when to take your first dose. Write it down to help you remember.

Your DOPTelet schedule:

DAY 1: _____

DAY 4: _____

DAY 2: _____

DAY 5: _____

DAY 3: _____

YOUR PROCEDURE DATE: _____

If you have any questions about how or when to take DOPTelet, call your **healthcare provider**.

You can also call your specialty pharmacy, or Dova 1Source™ at **1-833-DOVA-ONE (1-833-368-2663)** and they will help you.

Please see full Important Safety Information on page 11 and accompanying Full Prescribing Information for DOPTelet (avatrombopag), including Patient Information; also available at Doptelet.com.



DOPTelet is a registered trademark of AkaRx, Inc.
©2019 Dova Pharmaceuticals, Inc.
All rights reserved. PM-US-DOP-0158 Printed in USA.

Doptelet®
(avatrombopag) tablets