

HOW TO TAKE DOPTELET[®]

**Dosing to treat
low blood platelet counts
in adults with chronic immune
thrombocytopenia (cITP)
when other treatments
have not worked well enough.**

IMPORTANT SAFETY INFORMATION FOR DOPTELET

Before you take DOPTELET, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had a blood clot.
- are pregnant or plan to become pregnant. DOPTELET may harm your unborn baby. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with DOPTELET.
- are breastfeeding or plan to breastfeed. It is not known if DOPTELET passes into your breast milk. Do not breastfeed during your treatment with DOPTELET and for at least 2 weeks after the last dose. Talk to your healthcare provider about the best way to feed your baby during this time.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. DOPTELET may affect the way other medicines work, and other medicines may affect the way DOPTELET works.

**Please see full Important Safety Information on pages 4-5.
Full Prescribing Information in pocket.**

How should I take Doptelet?

Doptelet is a pill that is taken by mouth. Your healthcare provider will tell you how much Doptelet to take and when to start taking it.

Three things you should know about taking Doptelet:



1. Take exactly as instructed

Doses vary from person to person. Your healthcare provider will provide your dosing schedule.



2. Take with any type of food

Always remember to take Doptelet with food on a consistent schedule.



3. Your dose may change

Your doctor will monitor your platelet count to see if another dose may improve results.

Other things to remember when taking Doptelet:

- If you miss a dose of Doptelet, take it as soon as you remember and take your next dose at your usual scheduled time
- Do not take 2 doses at one time to make up for a missed dose. If you take too much Doptelet, call your healthcare provider or go to the nearest hospital emergency room right away

R_x Patient Name: _____
Address: _____

Prescription:
*Doptelet® (avatrombopag)
20 mg oral tablet
Dispense 30-day supply*

Directions:
*Take one 20 mg tablet
by mouth per day (QD)
with food*

Note: Example prescription depicts a starting dose.



IMPORTANT SAFETY INFORMATION FOR DOPELET (cont'd)

What are the possible side effects of DOPELET?

DOPELET may cause serious side effects, including blood clots. People with chronic immune thrombocytopenia and people with certain blood clotting conditions may have an increased risk of developing blood clots. Tell your healthcare provider right away if you have signs and symptoms of a blood clot, including:

- swelling, pain, or tenderness in your legs
- fast heartbeat
- shortness of breath
- stomach (abdominal) pain or tenderness
- chest pain

Please see full Important Safety Information on pages 4-5. Full Prescribing Information in pocket.

INDICATION AND IMPORTANT SAFETY INFORMATION

What is DOPTelet[®] (avatrombopag) [dop-TELL-it]?

DOPTelet is a prescription medicine used to treat low blood platelet counts in adults with chronic immune thrombocytopenia (ITP) when other treatments have not worked well enough.

DOPTelet is not used to make platelet counts normal in adults with chronic immune thrombocytopenia. It is not known if DOPTelet is safe and effective in children.

IMPORTANT SAFETY INFORMATION FOR DOPTelet

Before you take DOPTelet, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had a blood clot.
- are pregnant or plan to become pregnant. DOPTelet may harm your unborn baby. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with DOPTelet.
- are breastfeeding or plan to breastfeed. It is not known if DOPTelet passes into your breast milk. Do not breastfeed during your treatment with DOPTelet and for at least 2 weeks after the last dose. Talk to your healthcare provider about the best way to feed your baby during this time.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. DOPTelet may affect the way other medicines work, and other medicines may affect the way DOPTelet works.

How should I take DOPTelet?

- Take DOPTelet exactly as your healthcare provider tells you to take it.
- Your healthcare provider will tell you how much DOPTelet to take and when to start taking it.
- Your healthcare provider may change your dose of DOPTelet depending on your blood platelet counts.
- Take DOPTelet with food.
- If you take DOPTelet to treat your low blood platelet counts due to chronic immune thrombocytopenia, your healthcare provider will check your platelet count before, during and for at least 4 weeks after stopping your treatment with DOPTelet.
- If you are taking DOPTelet for chronic immune thrombocytopenia and you miss a dose of DOPTelet, take it as soon as you remember. Do not take 2 doses at one time to make up for a missed dose. Take your next dose at your usual scheduled time.
- If you take too much DOPTelet, call your healthcare provider or go to the nearest hospital emergency room right away.

What are the possible side effects of DOPTelet?

DOPTelet may cause serious side effects, including blood clots. People with chronic immune thrombocytopenia and people with certain blood clotting conditions may have an increased risk of developing blood clots. Tell your healthcare provider right away if you have signs and symptoms of a blood clot, including:

- swelling, pain, or tenderness in your legs
- shortness of breath
- chest pain
- fast heartbeat
- stomach (abdominal) pain or tenderness

The most common side effects of DOPTelet when used to treat low blood platelet counts in adults with chronic immune thrombocytopenia (ITP) are:

- headache
- tiredness
- bruising
- nosebleed
- upper respiratory tract infection
- joint pain
- bleeding gums
- purple or red spots on your skin
- runny nose

These are not all the possible side effects of DOPTelet.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store DOPTelet?

- Store DOPTelet at room temperature between 68°F to 77°F (20°C to 25°C).
- Store DOPTelet tablets in the original package.

Keep DOPTelet and all medicines out of the reach of children.

General information about the safe and effective use of DOPTelet.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use DOPTelet for a condition for which it was not prescribed. Do not give DOPTelet to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about DOPTelet that is written for health professionals.

What are the ingredients in DOPTelet?

Active ingredient: avatrombopag

Inactive ingredients: lactose monohydrate, colloidal silicon dioxide, crospovidone, magnesium stearate and microcrystalline cellulose. Tablet coating film: polyvinyl alcohol, talc, polyethylene glycol, titanium dioxide and ferric oxide yellow.

Please see accompanying full Prescribing Information, including the Patient Information Leaflet in pocket.

This Patient Information has been approved by the U.S. Food and Drug Administration.

WHAT IS DOPTELET CONNECT?

Doptelet®
connect

Doptelet Connect has Dedicated Case Managers that can help:

-  **Evaluate** a patient's prescription coverage, including benefits investigation, prior authorization, claims, and appeal assistance support
-  **Provide** field reimbursement support through a single point of contact
-  **Identify** potential financial assistance options that may be available to help patients with financial needs

For more information call 1-833-368-2663.

Case Managers are available Monday-Friday, 8 AM-8 PM ET.

DISCOVER THE DOPTELET COPAY ASSISTANCE PROGRAM

Eligible commercially insured patients may qualify for the Doptelet Copay Assistance Program

What's the program?

- Eligible patients may pay as little as \$0 for each Doptelet prescription
- Patients may save up to a maximum of \$15,000 per calendar year

Eligible patients must:

- Have commercial insurance that covers Doptelet
- Not be enrolled in any state or federal healthcare program such as Medicare, Medicaid, Medigap, Veterans Affairs (VA), Department of Defense (DOD), or TRICARE
- Be 18 years of age or older
- Be a resident in the United States or a US Territory

How to enroll:

- Approved pharmacies, healthcare professionals, and Doptelet Connect can enroll patients
- Patients can enroll themselves or can be enrolled by their caregivers
- Please call 1-833-368-2663 Monday-Friday, 8 AM-8 PM ET for more information



Doptelet
(avatrombopag) tablets

 **sobi**
rare strength

Doptelet is a registered trademark of AkaRx, Inc., a Sobi company.
Sobi is a trademark of Swedish Orphan Biovitrum AB (publ)
©2023 Sobi, Inc. - All rights reserved. PP-18437 06/23 Printed in the USA.