


COULD THIS CHRONIC ITP TREATMENT BE RIGHT FOR YOUR PATIENTS?

Doptelet is the only oral TPO-RA that can be taken anywhere, with any food on a consistent schedule.^{1,2*}



PATIENT CASE COURTESY OF DR TERRY GERNSHEIMER

BASED ON A REAL PATIENT†



CHRONIC ITP PATIENT: VIRGINIA

27-YEAR-OLD FEMALE

Treated with Doptelet after being nonresponsive to multiple courses of steroids

- Presenting symptoms: fatigue, bruising, and menorrhagia
- Pre-ITP diagnosis treatment: oral iron treatment and oral contraceptive pills
- Treatment after diagnosis: prescribed multiple corticosteroids and experienced adverse effects. Transitioned off corticosteroids to Doptelet
- At Doptelet initiation: 20 mg of Doptelet and 30 mg of prednisone
- Concomitant treatment reduction: discontinued prednisone and iron treatment after a month on Doptelet
- Maintenance platelet count on Doptelet: between $\geq 50 \times 10^9/L$ and $\leq 100 \times 10^9/L$

In a core study with Doptelet, patients achieved platelet goals of $50,000/\mu L$ (primary endpoint). Patients on Doptelet reached target platelet counts of $50,000/\mu L$ for a median of 12.4 cumulative weeks (primary endpoint).[‡]

"The best option for her was something that wouldn't be interfering with her diet or with her iron pills. And so we decided to go with Doptelet." — DR GERNSHEIMER^{1,3}

CLICK TO SEE DR GERNSHEIMER DISCUSS THIS PATIENT CASE

ITP=immune thrombocytopenia; TPO-RA=thrombopoietin receptor agonists.


* Food required.¹

† Photo is not the actual patient.

‡ Efficacy was evaluated in a 6-month, multicenter, randomized, double-blind, placebo-controlled Phase 3 study. Patients had received one or more prior chronic ITP therapies and had average screening and baseline platelet counts of $< 30 \times 10^9/L$. Forty-nine patients were randomized (2:1) to receive either Doptelet (n=32) or placebo (n=17).¹

DOPTELET PATIENT CASE TITRATION SCHEDULE¹

Platelets over time following Doptelet initiation



PLEASE SEE FULL IMPORTANT SAFETY INFORMATION CONTINUED ON NEXT PAGE AND FULL PRESCRIBING INFORMATION FOR DOPTELET AT WWW.DOPTELETHCP.COM.

INDICATION

DOPTELET is indicated for the treatment of thrombocytopenia in adult patients with chronic immune thrombocytopenia who have had an insufficient response to a previous treatment.

IMPORTANT SAFETY INFORMATION WARNINGS AND PRECAUTIONS

Thrombotic/Thromboembolic Complications. DOPTELET is a thrombopoietin (TPO) receptor agonist and TPO receptor agonists have been associated with thrombotic complications in patients with chronic liver disease (0.4%; (1/274) in DOPTELET-treated patients) and thromboembolic complications in patients with chronic immune thrombocytopenia (7%; (9/128) in DOPTELET-treated patients). Portal vein thrombosis has been reported in patients with chronic liver disease, and thromboembolic events (arterial and venous) have been reported in patients with chronic immune thrombocytopenia treated with TPO receptor agonists.

Doptelet
(avatrombopag) tablets



BROAD INSURANCE COVERAGE

Doptelet is available for 97% of patients with commercial insurance and the majority of Medicare-insured patients nationwide^{4,5*}



SEE THE FORMULARY STATUS IN YOUR AREA
www.DopteletCoverage.com

Available resources include:

- Prior Authorization Guide
- Letter of Medical Necessity (Fillable)
- Letter of Appeal (Fillable)



Request FRM
A Field Reimbursement Manager (FRM) can educate on access processes
www.doptelethcp.com

Help your patients access the Doptelet lift



Discover patient support programs for Doptelet
www.doptelethcp.com/itp/access

*Majority is defined as >50% of patients covered.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Consider the potential increased thrombotic risk when administering DOPTelet to patients with known risk factors for thromboembolism, including genetic prothrombotic conditions and acquired risk factors.

DOPTelet should not be administered to patients with chronic liver disease or chronic immune thrombocytopenia in an attempt to normalize platelet counts. Monitor platelet counts, and for signs and symptoms of thromboembolic events and institute treatment promptly.

Serious Adverse Reactions

Serious adverse reaction that occurred more frequently in patients treated with DOPTelet (9%; 12/128) compared to placebo (5%; 1/22) was headache, occurring in 1.6% (2/128).

Adverse Reactions

The most common adverse reactions ($\geq 10\%$) in patients with chronic immune thrombocytopenia were headache, fatigue, contusion, epistaxis, upper respiratory tract infection, arthralgia, gingival bleeding, petechiae, and nasopharyngitis.

References: **1.** DOPTelet (avatrombopag) [prescribing information]. Durham, NC: AkaRx, Inc; 2021. **2.** Promacta (eltrombopag) [prescribing information]. East Hanover, NJ: Novartis Pharmaceuticals Corp; 2023. **3.** McDonald V, Newland A, Morgan M, et al. Patient preferences and experiences regarding thrombopoietin-receptor agonists for immune thrombocytopenia in the United Kingdom and Ireland (TRAPeze UK & IE study). *Hematology*. 2021;26(1): 799-808. **4.** Data on file. MMIT commercial summary export. 2024: Sobi, Inc. **5.** MMIT Medicare summary export 2024: Sobi, Inc.

Postmarketing Experience

Following the approval of DOPTelet, hypersensitivity reactions involving the immune system, including, but not limited to, pruritus, rash, choking sensation, swollen face, and swollen tongue have been reported.

These are not all the possible risks associated with DOPTelet. Please see Full Prescribing Information for DOPTelet at www.doptelethcp.com

To report suspected adverse reactions, contact Sobi North America at 1-866-773-5274 or FDA at 1-800-FDA-1088.

For statutory pricing disclosures, visit doptelethcp.com/wac-pricing.



Doptelet is a registered trademark of AkaRx, Inc., a Sobi company.
Sobi is a registered trademark of Swedish Orphan Biovitrum AB (publ)
© 2025 Sobi, Inc. - All rights reserved.
PP-22354v3 02/25

Doptelet
(avatrombopag) tablets